


Cosmological constraints from Chandra observations of galaxy clusters



Steve Allen, IoA, Cambridge. UK.

With R. Schmidt (Potsdam)
A. Fabian (Cambridge)
S. Bridle (Cambridge)
H. Ebeling (Hawaii)
L. van Speybroeck (CfA)
A. Edge (Durham)

PROJECT OUTLINE:

Last four years, we have carried out a programme to study most X-ray luminous, dynamically relaxed galaxy clusters identified from ROSAT All-Sky Survey.

Aims: measure mass distributions using Chandra X-ray data and independent gravitational lensing studies \rightarrow improved constraints on cosmological parameters ($\Omega_m, \Omega_\Lambda, \sigma_8, w \dots$)

THIS TALK (cosmological constraints):

Constraints on Ω_m, Ω_Λ from the X-ray gas mass fraction, f_{gas} , in dynamically relaxed clusters and its apparent redshift dependence.

- precise constraint on Ω_m .
- confirmation of SNIa results on dark energy.

Constraints on σ_8, Ω_m from local X-ray luminosity function (XLF).

Results from combination of Cosmic Microwave Background (WMAPext) and Chandra $f_{\text{gas}}(z)$ data.

- combination of data sets breaks degeneracies in each
 - \rightarrow tight constraints on key cosmo. params. (H_0, Ω_q, w).
- (with XLF data) \rightarrow preference for a non-zero neutrino mass.

X-RAY MASS MEASUREMENTS:

Divide clusters into circular annuli (2D) or spherical shells (3D).

- 1) observed X-ray surface brightness profile
2) deprojected (spectrally-determined) kT profile

+ assump. hydrostatic equilibrium (spherical symmetry) → $M(r)$.

In practise, studies of distant clusters require modified approach:
Take simple parameterized mass model (NFW: r_s, c) + SB(r) →
run Monte-Carlo simulations predict $kT(r)$ → compare with obs.

Examine grid 100×100 parameter values for each mass model.
Evaluate χ^2 for each grid point → best-fit model + uncertainties.

LENSING STUDIES:

Weak lensing: ($r \gtrsim 200h_{50}^{-1}$ kpc)



Results from literature + own programmes underway.

Strong lensing: ($r \lesssim 200h_{50}^{-1}$ kpc)



Ground based programme complete. HST (WFPC2/ACS) programme underway.

KEY: Combination of independent X-ray plus lensing
methods → robust results!

X-RAY + WEAK LENSING:



STRONG LENSING IN MS2137.3-2353:



Excellent agreement between observed and predicted arc configurations, only free parameters background source positions.

Cosmological constraints from the X-ray gas mass fraction in the most luminous, relaxed galaxy clusters

References: White & Frenk (1991). Fabian (1991), White *et al.* (1993), David *et al.* (1995), White & Fabian (1995), Evrard (1997), Ettori & Fabian (1999), Allen *et al.* (2002, 2003), Ettori *et al.* (2003), Arnaud *et al.* (2003).

BASIC IDEA: Galaxy clusters are so large that their matter content should provide a fair sample of matter content of Universe.

Chandra + lensing data \rightarrow robust total mass measurements

Chandra data \rightarrow (very) precise X-ray gas mass measurements

X-ray gas dominates baryonic mass content of clusters (6-7 times more mass than all stars in galaxies *e.g.* Fukugita *et al.* 1998).

If we define:


$$f_{\text{gas}} = \frac{\text{Xray gas mass}}{\text{total mass in cluster}} \quad f_{\text{gal}} = 0.19h^{0.5} f_{\text{gas}}$$

$$\text{Then} \quad f_{\text{baryon}} = f_{\text{gal}} + f_{\text{gas}} = f_{\text{gas}}(1 + 0.19h^{0.5}).$$

Since clusters provide fair sample of Universe, $f_{\text{baryon}} = \Omega_{\text{b}}/\Omega_{\text{m}}$.

$$\Omega_{\text{m}} = \frac{\Omega_{\text{b}}}{f_{\text{baryon}}} = \frac{\Omega_{\text{b}}}{f_{\text{gas}}(1 + 0.19h^{0.5})}$$

So given f_{gas} , Ω_{b} (cosmic nucleosynthesis, CMB) $\rightarrow \Omega_{\text{m}}$.



(fig from Allen *et al.* 2003. Data analysed here has 10 clusters.)

$f_{\text{gas}}(r) \rightarrow$ approximately universal value at $r \sim r_{2500}$

$$\begin{aligned} \rightarrow \text{weighted mean } \bar{f}_{\text{gas}}(r_{2500}) &= (0.110 \pm 0.003)h_{70}^{-1.5} \\ &= (0.065 \pm 0.002)h^{-1.5} \end{aligned}$$


$$\begin{aligned} \text{Given } \Omega_b h^2 &= 0.0214 \pm 0.0020 && (\text{Kirkman } et al. \ 2003) \\ h = H_0/100 &= 0.72 \pm 0.08 && (\text{Freedman } et al. \ 2001) \end{aligned}$$

$$\begin{aligned} \Rightarrow \Omega_m &= \frac{(0.0214 \pm 0.0020)h^{-0.5}}{(0.065 \pm 0.002)(1 + 0.19 h^{0.5})} \\ &= 0.323 \pm 0.031 \end{aligned}$$

Indication from simulations that baryonic mass fraction in clusters is slightly lower than mean value for universe as a whole.

$$f_{\text{baryon}} = b \frac{\Omega_{\text{b}}}{\Omega_{\text{m}}}$$

e.g. Eke, Navarro & Frenk (1998)



For $r = 0.25 r_{\text{vir}}$ (Chandra obs.) $b = 0.83 \pm 0.04$


$$\longrightarrow \Omega_{\text{m}} = 0.27 \pm 0.03$$

Apparent variation of f_{gas} with redshift:

When measuring f_{gas} we adopt a reference cosmology. Since $f_{\text{gas}} \propto D_A^{1.5}$, this introduces apparent, systematic redshift variations, depending on differences between adopted and underlying cosmology.

If clusters provide a fair sample of matter content of the Universe then we expect $f_{\text{gas}}(z)$ - measured within r_{2500} - to be constant.

When adopted cosmology='true' cosmology, expect $f_{\text{gas}}(z) = \text{const.}$




Can easily see $f_{\text{gas}}(z)$ data favour Λ CDM over SCDM and that standard Λ CDM ($\Omega_m = 0.3, \Omega_\Lambda = 0.7$) close to 'true' cosmology.

Above figures are for older data of Allen *et al.* 2003. Left SCDM. Right Λ CDM (0.3, 0.7). Current data used in analysis reported here contains 10 clusters with $z < 0.6$. (Allen *et al.* , in preparation.)

To quantify: fit SCDM data with model which accounts for expected apparent variation in $f_{\text{gas}}(z)$ data as underlying cosmology is varied (Ω_m, Ω_Λ) \rightarrow find cosmology giving best-fit to data.

$$f_{\text{gas}}^{\text{mod}}(z) = \frac{b \Omega_b}{(1 + 0.19\sqrt{h}) \Omega_m} \left[\frac{h}{0.5} \frac{D_A^{\Omega_m=1, \Omega_\Lambda=0}(z)}{D_A^{\Omega_m, \Omega_\Lambda}(z)} \right]^{1.5}$$

$$(\Omega_b h^2 = 0.0214 \pm 0.0020, \quad h = 0.72 \pm 0.08, \quad b = 0.83 \pm 0.04)$$




\Rightarrow Best fit: $\Omega_m = 0.267 \pm 0.034, \quad \Omega_\Lambda = 0.78 \pm 0.33$

Good fit: reduced $\chi^2 \sim 0.6$

(Above figure is for older data. Current data used in analysis here contains 10 clusters. Allen *et al.* , in prep.)

Marginalized results on dark energy (Λ CDM)



Red curve: standard priors


Blue curve: weak priors $h=0.72 \pm 0.20$, $b = 0.83 \pm 0.1$,
 $\Omega_b h^2 = 0.0214 \pm 0.006$

First direct confirmation of SN1a results, revealing effects of dark energy on a single source population as a function of redshift.

Future Chandra/XMM observations could improve significance of detection to $> 4\sigma$ over next few yrs (targets identified) allowing us to probe dw/dz and test nature of dark energy.

(Allen *et al.* , in prep.)

Comparison with independent constraints Ω_m, Ω_Λ



Chandra $f_{\text{gas}}(z)$ data. 1, 2 and 3σ contours.

CMB (COBE + Boomerang'98 + MAXIMA-1; Jaffe *et al.* 2001).

Supernovae (Riess *et al.* 1998; Perlmutter *et al.* 1999).

+ also consistent with 2dF + CMB (*e.g.* Efsthathiou *et al.* 2002, Lahav *et al.* 2002; Percival *et al.* 2002, Lewis & Bridle 2002).

Agreement between independent analyses/methods.

All results consistent within 1σ statistical uncertainties

$$\Omega_m \sim 0.3, \quad \Omega_\Lambda \sim 0.7$$

Constraints on Ω_m and σ_8 from the local X-ray luminosity function of the most X-ray luminous galaxy clusters.

References: *e.g.* Evrard 1989, Henry & Arnaud 1991, Henry *et al.* 1992, Oukbir & Blanchard 1992, White *et al.* 1993, Eke *et al.* 1996, Viana & Liddle 1996, Kitayama & Suto 1997, Borgani *et al.* 1997, Markevitch 1998


THIS ANALYSIS (Allen, Schmidt, Fabian, Ebeling 2003):

- 1) Local ($z < 0.3$) X-ray luminosity function (XLF) of clusters from ROSAT All-Sky Survey (RASS): eBCS (north) and REFLEX (south) studies (Ebeling *et al.* 2000; Böhringer *et al.* 2002).
- 2) Mass-luminosity relation from Chandra and ROSAT X-ray observations + weak lensing measurements
→ Precise (M_{200}) mass, luminosity measurements for 17 clusters (including both relaxed and dynamically active systems).
- 3) Predicted mass function of clusters from Hubble Volume simulations for Λ CDM (Jenkins *et al.* 2001; Evrard *et al.* 2002).
- 4) Chandra $f_{\text{gas}}(z)$ data for dynamically relaxed clusters.

combine → constraints on σ_8 and Ω_m .

The observed local XLF of galaxy clusters

Compilation of recently published samples (Rosati *et al.* 2002):




Excellent agreement between independent studies based on ROSAT All-Sky Survey and pointed ROSAT/Einstein observations.

Here, work with XLF from RASS: eBCS (north) and REFLEX (south) studies (Ebeling *et al.* 2000; Böhringer *et al.* 2002).

→ 111 clusters with $L_{X,0.1-2.4} > 10^{45} \text{ erg s}^{-1}$ ($h = 0.5 \Lambda \text{CDM}$)

Luminosity cut keeps systematic uncertainties to a minimum. These are clusters for which virial relations are calibrated most precisely.

The observed mass-luminosity relation ($\Delta = 200$)



Masses, M_{200} , measured from Chandra and weak lensing data.

0.1 – 2.4 keV luminosities from pointed ROSAT observations.

Evolution parameter $E(z) = (1+z)\sqrt{(1+z\Omega_m + \Omega_{\Lambda}/(1+z)^2 - \Omega_{\Lambda})}$
(assume Λ CDM, $h = 0.5$ cosmology to match BCS/REFLEX).

Characterize data using simple power-law model


$$E(z) \left[\frac{M_{200}}{1 M_{\odot}} \right] = M_0 \left[\frac{L}{E(z) 10^{44} \text{ erg s}^{-1}} \right]^{\alpha}.$$

$$(M_0 = 2.0 \pm 1.0 \times 10^{14} M_{\odot}, \alpha = 0.76 \pm 0.15)$$

Construct model XLF

Combine observed mass-luminosity relation + predicted mass function (Evrard *et al.* 2002) \rightarrow predicted model XLF (σ_8 , Ω_m).

Fit model XLF to observed XLF \rightarrow constraints on σ_8 , Ω_m .




Simple χ^2 approach:

Using best-fit M-L relation, we find that model with $\sigma_8 = 0.73$, $\Omega_m = 0.25$ provides best fit to the observed XLF ($\chi^2_{\min} = 5.1/4$).

Results on σ_8 as a function of Ω_m

Use Monte Carlo analysis which accounts for effects of scatter as well as uncertainties in norm and slope of M-L relation.

Flat Λ CDM cosmology:




$$\sigma_8 = (0.508 \pm 0.019) \Omega_m^{-(0.253 \pm 0.024)}$$

$$\Omega_m < 0.34$$

Comparison with independent results (flat Λ CDM)

Other studies based on local cluster abundance



Seljak (2002). Local XTF, observed M-T

Reiprich & Böhringer (2001). Local XTF, observed M-T

Viana *et al.* (2002). Local XLF . Stacked lensing M-L


Bahcall *et al.* (2002). Optical (SDSS) clusters, mass-richness

Consistent with local XLF, XTF analyses of Schuecker *et al.* (2003), Pierpaoli *et al.* (2003) + study of local baryonic mass function by Voevodkin & Vikhlinin (2003).

Breaking the $\sigma_8 - \Omega_m$ degeneracy

The degeneracy between σ_8 and Ω_m from local XLF/XTF studies can be broken by combining with Chandra $f_{\text{gas}}(z)$ data.

Include Gaussian priors: $\Omega_b h^2 = 0.0205 \pm 0.0018$, $h = 0.72 \pm 0.08$
 $b = 0.83 \pm 0.04$, $\Omega_k = 0.0$.




→ Best-fit and marginalized 68 per cent confidence limits:

$$\sigma_8 = 0.71 \pm 0.04$$

$$\Omega_m = 0.27 \pm 0.03$$

Cosmological constraints from CMB+ $f_{\text{gas}}(z)$ data.



Extended WMAP
data set

WMAP (TT/TE)
+CBI+ACBAR

Analyse CMB (WMAP+CBI+ACBAR) data using MCMC method.
Importance sample the results folding in $f_{\text{gas}}(z)$ constraints.


Use 8 parameter model with free parameters

$\Omega_{\text{dm}}h^2$	physical dark matter density
$\Omega_{\text{b}}h^2$	physical baryonic matter density
H_0	Hubble constant
A_{S}	scalar amplitude
n_{S}	scalar spectral index
z_{rec}	redshift of recombination
Ω_{k}	spatial curvature
w	quintessence parameter = p/ρ
R	tensor/scalar amplitude ratio
n_{T}	tensor spectral index
f_{ν}	neutrino mass fraction

Combination of CMB+ $f_{\text{gas}}(z)$ data breaks degeneracies present in both data sets

$$f_{\text{gas}}^{\text{mod}}(z) \propto \frac{\Omega_{\text{b}}}{\Omega_{\text{m}}} \left[\frac{h}{0.5} \frac{D_{\text{A}}^{\Omega_{\text{m}}=1, \Omega_{\text{q}}=0}(z)}{D_{\text{A}}^{\Omega_{\text{m}}, \Omega_{\text{q}}}(z)} \right]^{1.5}$$


e.g. constraints on Ω_{m} , H_0 (Ω_{k} free, w free)



Black: CMB only Blue: CMB+ $f_{\text{gas}}(z)$

$$\Omega_{\text{m}} = 0.29^{+0.04}_{-0.03}, \quad H_0 = 68 \pm 6 \text{ km s}^{-1} \text{ Mpc}^{-1}$$

Dark energy density: quintessence models (Ω_k, w free)



Black: CMB only


Blue: CMB + $f_{\text{gas}}(z)$

Marginalized results:

$$\Omega_m = 0.29^{+0.04}_{-0.03}, \quad \Omega_q = 0.73^{+0.03}_{-0.04}$$

(Allen *et al.* , in prep.)

Quintessence parameter ($w \equiv p/\rho$)



Marginalized results:


$$\begin{aligned} f_{\text{gas}}(z)+\text{CMB } (\Omega_{\text{k}} \text{ free}) : w &= -1.17 \pm 0.26 \\ f_{\text{gas}}(z)+\text{CMB } (\text{flat}) : w &= -1.02^{+0.17}_{-0.31} \\ f_{\text{gas}}(z)+\text{BBNS}+\text{HST } (\text{flat}) : w &= -1.04^{+0.31}_{-0.36} \end{aligned}$$

→ results consistent with $w = -1$ (Λ CDM)

(Allen *et al.* , in prep.)


Preference for a non-zero neutrino mass

CMB+2dF+ $f_{\text{gas}}(z)$ data: Flat geometry, $w=-1$, allow tensors.



Assuming negligible
neutrino mass

$$f_\nu = \Omega_\nu / \Omega_m = 0$$




With neutrino mass
as free parameter

$$\rightarrow \Omega_\nu / \Omega_m = 0.05$$

(Allen, Schmidt & Bridle 2003)

Marginalized constraints on neutrino density



Tentative detection of non-zero species-summed neutrino mass

$$\sum_i m_i \sim 94 \text{ eV } \Omega_\nu h^2 = 0.56^{+0.30}_{-0.26} \text{ eV}$$

Given small mass-squared differences from solar and atmospheric neutrino experiments, and assuming 3 active neutrino species (SLAC, LEP) and an absence of massive, sterile neutrinos

→ Approximately degenerate neutrino mass.

$$m_\nu = 0.19^{+0.10}_{-0.09} \text{ eV} \quad (\text{CMB}+2\text{dF}+f_{\text{gas}}+\text{XLF})$$

$$m_\nu = 0.17^{+0.22}_{-0.05} \text{ eV} \quad (\text{CMB}+f_{\text{gas}}+\text{XLF})$$

Consistent with 2dF team ($\sum_i m_i < 2.2 \text{ eV}$) and marginally with WMAP-team, including Ly- α forest data ($\sum_i m_i < 0.71 \text{ eV}$)

CONCLUSIONS

Chandra + grav. lensing data → precise mass measurements for dynamically relaxed clusters. Mass profiles well described by NFW models with parameters (r_s, c) consistent with simulations.

Chandra $f_{\text{gas}}(z)$ data for relaxed clusters in combination with other data → tight constraint on mean matter density of Universe, Ω_m .

$$f_{\text{gas}}(z) + \text{BBN} + \text{HST} \longrightarrow \Omega_m = 0.27 \pm 0.03$$

$$f_{\text{gas}}(z) + \text{CMB} \longrightarrow \Omega_m = 0.29^{+0.04}_{-0.03}$$

Chandra $f_{\text{gas}}(z)$ data → direct confirmation of SN1a results on DE and interesting constraints on quintessence.

$$f_{\text{gas}}(z) + \text{weak priors} \longrightarrow \Omega_\Lambda = 0.78 \pm 0.33$$

$$f_{\text{gas}}(z) + \text{CMB} \longrightarrow \Omega_q = 0.73^{+0.03}_{-0.04}$$

$$\longrightarrow w = -1.17 \pm 0.26$$

Local XLF + observed M/L → tight constraint on amplitude of mass fluctuations on $8 h^{-1}\text{Mpc}$ scales, $\sigma_8 \sim 0.7$ for $\Omega_m \sim 0.3$.

Combination of CMB + 2dF + $f_{\text{gas}}(z)$ + XLF data → preference for a non-zero neutrino mass → $m_\nu = 0.19^{+0.10}_{-0.09} \text{ eV}$.